

search

Brood parasitism in fish

May 10, 2018, University of Konstanz



Adult cuckoo catfish (*synodontis multipunctatus*). Credit: Institute of Vertebrate Biology, Brno (Czech Republic)

There are other animals besides the cuckoo who smuggle their offspring into another animal's nest. The *synodontis multipunctatus*, which lives in Lake Tanganyika in Africa and is better known as the cuckoo catfish, is just as cunning as the cuckoo. This savvy parasite, like the cuckoo bird, places its eggs among those of cichlids. To protect their eggs, cichlids carry their eggs in their mouths.

Professor Axel Meyer, an evolutionary biologist from the University of Konstanz, and a team of researchers from the Institute of Vertebrate Biology in Brno (Czech Republic) have carried out research into the evolutionary strategies employed by cuckoo catfish and various types of cichlids that occur in Lake Tanganyika and several other African lakes. Their study paints a fascinating picture of evolutionarily shaped and individually learned defence behaviour, as well as the deception efforts

employed by both species of fish—and the high price that cichlids pay for keeping the illegitimate offspring of the cuckoo catfish away from their own eggs.

The research findings were published in the *Science Advances* issue published on 2 May 2018.

Lake Tanganyika in Africa is famous for its biodiversity. Many of its 250 endemic species of cichlids are mouthbreeders. To protect their offspring and prevent other fish from devouring them, cichlids carry and breed their eggs in

Featured	Last comments	Popular
	May 07, 2018	13
	May 07, 2018	62
	May 07, 2018	10
	May 07, 2018	2
	May 07, 2018	0

[more »](#)

Phys.org on Facebook

Email newsletter

their mouths. For several weeks after hatching and swimming by themselves, the young fish return to their mother's mouth for protection.

It is this very particular brood care behaviour that the cuckoo catfish, also endemic to Lake Tanganyika, has learned to exploit. When the cichlids spawn, the cuckoo catfish places its own eggs among a cichlid's clutch of eggs. If this goes unnoticed by the cichlid, and if it distinguish its own eggs from those of the catfish, it will carry and breed both its own and the catfish eggs in its mouth. However, the larvae of the cuckoo catfish hatch sooner, devouring the cichlid's own offspring. Often, the cichlid will believe the illegitimate offspring of the catfish to be her own, continuing to protect them.

But the cichlids have learned to defend themselves against the cunning of the cuckoo catfish. When gathering their eggs into their mouth, they try to identify and exclude the smuggled eggs. Often, however, overcaution will lead them to reject some of their own eggs, as well—a high price in return for their own evolutionary fitness.



Cichlid (*simochromis diagramma*) from Lake Tanganyika. Credit: Institute of Vertebrate Biology, Brno (Czech Republic)

"Both species of fish have co-evolved for millions of years," says Axel Meyer about their well-matched relationship of deception and defence. The behaviour of these two species is evidence of what the biologist calls "evolutionary experience," which he documented in his joint study with his colleagues from Brno.

Tableau - Official Site - Free

Answer Questions as Fast as You Can Think of
tableau.com

The scientists obtained eggs both from the cuckoo catfish and from the mouthbreeding cichlids that live in Lake Tanganyika and raised them in an aquarium. Then they compared the captive cichlids' capacity for distinguishing their own eggs with that of other types of cichlid from other bodies of water where cuckoo catfish do not occur. The result: The deceptive strategy employed by the cuckoo catfish worked between three and eleven times better on the "evolutionarily naive" cichlids from other bodies of water. Due to their evolutionary experience, the cichlids from Lake Tanganyika, who share an evolutionary history with the cuckoo catfish, were much more successful in identifying and rejecting the parasite's eggs. By the term "evolutionary experience," the scientists mean natural selection in favour of the ability to discriminate smuggled eggs.

The study also revealed that cichlids lacking evolutionary experience are unable to learn to reject the eggs of the cuckoo catfish—in contrast to co-

Relevant PhysicsForums posts

How to measure pain objectively? 1 hour ago

Seeking a cheap wide mouthed jar capable of withstanding a vacuum May 09, 2018

Brain circuit for this behavior May 08, 2018

Query about Vitamin ##K_1## and ##K_2## May 08, 2018

What type of bacteria evolved into mitochondria? May 08, 2018

Neuroscience and math May 07, 2018

More from Biology and Medical

evolved cichlids that increase their chances to see through the cuckoo catfish trick. This ability to adapt made the cichlids from Lake Tanganyika much more successful when coping with brood parasites. These findings suggest that is not the combination of evolutionary experience with individual experience and the ability to learn that help cichlids discriminate between their own and foreign eggs.

Several bird species are known to practice brood parasitism, i.e. the smuggling of eggs into another bird's nest. Among fish, the cuckoo catfish is the only known obligate brood parasite. None of the other 40 catfish species endemic to Lake Tanganyika are known to behave like this.

Explore further: Russian cuckoo invasion spells trouble for Alaskan birds, study finds

More information: Radim Blažek et al. Success of cuckoo catfish brood parasitism reflects coevolutionary history and individual experience of their cichlid hosts, *Science Advances* (2018). DOI: [10.1126/sciadv.aar4380](https://doi.org/10.1126/sciadv.aar4380)

Journal reference: [Science Advances](#)

38 shares

Provided by: [University of Konstanz](#)

[feedback to editors](#)

31		Tweet					
Like			reddit				

Related Stories



Russian cuckoo invasion spells trouble for Alaskan birds, study finds

May 7, 2018

Common cuckoos and oriental cuckoos in eastern Russia appear to be expanding their breeding range into western Alaska, where songbirds are naive to the cuckoos' wily ways, researchers report. A new study suggests the North ...



The dark side of cichlid fish—from cannibal to caregiver

April 20, 2018

Cannibalism, the eating of conspecifics, has a rational background in the animal kingdom. It may serve as a source of energy-rich nutrition or to increase reproductive success. Some species do not even spare their own brood. ...



Egg colours make cuckoos masters of

Recommended for you



What gives bees their sweet tooth?

May

10, 2018

Scientists have discovered bees linger on a flower, emptying it of nectar, because they have sugar-sensing taste neurons which work together to prolong the pleasure of the sweetness.



Neuroscientists find first evidence animals can mentally replay past events

May 10, 2018

Neuroscientists at Indiana University have reported the first evidence that non-human animals can mentally replay past events from memory. The discovery could help advance the development of new drugs to treat Alzheimer's ...



Ancient skull shows early 'baleen whale'

User comments

Please sign in to add a comment. Registration is free, and takes less than a minute. [Read more](#)

[Click here to reset your password.](#)

[Sign in to get notified via email when new comments are made.](#)

We recommend

First insights into the diversity of gill monogeneans of 'Gnathochromis' and *Limnochromis* (Teleostei, Cichlidae) in Burundi: do the parasites mirror host ecology and phylogenetic history?

Nikol Kmentová et al., PeerJ

Using 3D printed eggs to examine the egg-rejection behaviour of wild birds.

Branislav Igic et al., PeerJ

Patterns and coevolutionary consequences of repeated brood parasitism

M. E. Hauber et al., Biology Letters

A gene expression study of dorso-ventrally restricted pigment pattern in adult fins of *Neolamprologus meeli*, an African cichlid species.

Ehsan Pashay Ahi et al., PeerJ

Chronic playback of boat noise does not impact hatching success or post-hatching larval growth and survival in a cichlid fish.

Rick Bruintjes et al., PeerJ

Extra-pair paternity in the long-tailed finch *Poephila acuticauda*.

Erica P van Rooij et al., PeerJ

Powered by

Top	Help	Science X Account	Feature Stories	Android app	Connect
Home	FAQ	Sponsored Account	Latest news	iOS app	
Search	About	Newsletter	Week's top	Amazon Kindle	
Mobile version	Contact	RSS feeds	Archive	Push notification	