

more than 261,000 arti

Search our Site:

[Home](#) [About us](#) [Deutsch](#)
[RSS-Feed](#)  
[SCIENCE REPORTS](#) [SPECIAL TOPICS](#) [B2B AREA](#) [JOBS & OPPORTUNITIES](#)

## REPORTS AND NEWS

[Agricultural and Forestry Science](#)  
[Architecture and Construction](#)  
[Automotive Engineering](#)  
[Business and Finance](#)  
[Communications Media](#)  
[Earth Sciences](#)  
[Ecology, The Environment and Conservation](#)  
[Health and Medicine](#)  
[Information Technology](#)  
[Interdisciplinary Research](#)  
[Life Sciences](#)  
[Machine Engineering](#)  
[Material Sciences](#)  
[Medical Engineering](#)  
[Physics and Astronomy](#)  
[Power and Electrical Engineering](#)  
[Process Engineering](#)  
[Social Sciences](#)  
[Studies and Analyses](#)  
[Transportation and Logistics](#)

[Home](#) [Science Reports](#) [Reports and News](#) [Life Sciences](#)

## Rapid evolution: New findings on its molecular mechanisms

**14.08.2019**

Evolutionary biologists from Konstanz analyze the role of microRNAs in the evolution of new species

The mechanisms by which new species arise are still not fully understood. What are the evolutionary processes that drive the evolution of new species? Evolutionary biologists traditionally assumed that geographical barriers between animal populations play a decisive role (allopatric speciation): a species is physically separated into two or more isolated populations, thereby preventing gene flow between these groups.

The subpopulations adapt to their respective habitats and evolve into independent species with different characteristics. In recent years, however, the evolutionary biologist Professor Axel Meyer from the University of Konstanz has not only been able to show that new species can evolve from a source population within a shared habitat and in the presence of gene flow (sympatric speciation), but that this type of speciation might be much more common than previously thought.



The Midas cichlid fish from the crater lakes of Nicaragua are one of the the best known examples for sympatric speciation. They evolved from a source

### Further sponsors


[VideoLinks](#)  
[Industry & Economy](#)



His laboratory is investigating both the ecological and genetic mechanisms that facilitate sympatric speciation. In a recent publication in the scientific journal *Molecular Biology and Evolution*, Axel Meyer and his colleagues Paolo Franchini, Peiwen Xiong, Carmelo Fruciano, Ralf Schneider, Joost Woltering and Darrin Hulsey identify the decisive role that a kind of genetic switch, microRNAs, play in sympatric speciation.

... more about:

» **Molecular Biology** » **cichlid fish** » **ecological niches** » **gene flow** » **genes** » **molecular mechanisms** » **new species** » **sympatric speciation**

Rapid evolution of cichlids

The researchers led by Axel Meyer analyze the Midas cichlid fish from Nicaragua's volcanic crater lakes as a model system in their research. Cichlids are known for their ability to adapt to new environments at an exceptionally rapid rate and to

form new species.

The new fish species originate from a population found in the large lakes of Nicaragua, but after colonizing several very small and young crater lakes, they adapted to new ecological niches, evolved new characteristics (e.g. a more elongated body or a different jaw) and evolved into a variety of new species in less than 22,000 years.

A further distinctive feature of these Midas cichlids is that they evolved into different species within the same population, sometimes repeatedly, by adapting to different ecological niches within their small crater lakes. The Midas cichlids of Nicaragua are thus one of the best known examples for sympatric speciation.

Within such a short time period new mutations are very unlikely to happen. This made finding a molecular mechanism that can bring about different body shapes, adaptations and thereby distinct ecological niches so difficult and interesting. The evolution of microRNAs and new target sites for them to regulate the expression of genes offers one potential molecular mechanism that can bring about very fast evolutionary change with very little genetic differences between the extremely young species.

microRNA

The biologists from Konstanz carried out genetic analyses of five species of Midas cichlids from the crater lakes Apoyo and Xiloá. Here, they focused in particular on the function of the so-called microRNA – a non-coding ribonucleic acid that has a regulatory effect on gene expression. The researchers found an increased activity of microRNA in young fish one day after they hatched during a phase in which the fish bodies are formed.

They analyzed the interaction between microRNA and gene expression, identifying specific pairs of microRNAs and genes that influence each other. The microRNA suppresses the expression of the target genes and thus has a regulatory effect on them: the more active a particular microRNA is, the more effectively the target gene is suppressed or "switched off". "Our research results provide strong evidence that extremely quickly evolving microRNA regulation contributes to the rapid sympatric speciation of Midas cichlids," says Paolo Franchini.

Facts:

- Evolutionary biologists from Konstanz investigate the role of microRNA in the rapid sympatric speciation of Midas cichlids – the evolution of new species within the same habitat and in the presence of gene flow.

- Original publication: Paolo Franchini, Peiwen Xiong, Carmelo Fruciano, Ralf F Schneider, Joost M Woltering, C Darrin Hulsey, Axel Meyer, MicroRNA gene regulation in extremely young and parallel adaptive radiations of crater lake cichlid fish, *Molecular Biology and Evolution*, msz168

Link: <https://academic.oup.com/mbe/advance-article/doi/10.1093/molbev/msz168/5545549>

- Genetic analyses of five species of Midas cichlids from the crater lakes Apoyo and Xiloá, which evolved out of a shared source population into new species in far less than 22,000 years.

- Analyses of the interaction between microRNA and genes. Specific pairs of microRNAs and genes that interact with each other were identified. The microRNA suppresses the expression of the target genes.

- This research project was carried out in the context of Professor Axel Meyer's ERC Advanced Grant "Comparative genomics of parallel evolution in repeated adaptive radiations" (GenAdap 293700). The project was also funded by a German Research Foundation (DFG) grant (FR 3399/1-1).

population into a variety of independent species in less than 22,000 years.

University of Konstanz



#### Event News

4th Hybrid Materials and Structures 2020 28 - 29 April 2020, Karlsruhe, Germany  
14.08.2019 | Event News

What will the digital city of the future look like? City Science Summit on 1st and 2nd October 2019 in Hamburg  
12.08.2019 | Event News

Jena Laser Technology Conference brings together top international researchers  
12.08.2019 | Event News

## find and help

to the campaign page >>>

#### Latest News

Artificial intelligence in the fight against river blindness  
14.08.2019 | Information Technology


Fracking prompts global spike in atmospheric methane  
14.08.2019 | Studies and Analyses

Rapid evolution: New findings on its molecular mechanisms  
14.08.2019 | Life Sciences

#### VideoLinks Science & Research



**Infrared emitters for the automotive industry**  
Re-activating of adhesives on automotive glass



**The incredible power of light!**  
Light is more than you can see.



DAIMLER

DEUTSCHE  
STEINZEUG

Q.CELLS

Roland Berger  
Strategy Consultants

PHILIPS

Heraeus

BBK

Allianz

3M



SAMSUNG

Deutsche Bank

businessAD

n-tv

BERTELSMANN  
media worldwideLufthansa Cargo  
The business to business class.

Note to editors:

You can download pictures here:

[https://cms.uni-konstanz.de/fileadmin/pi/fileserver/2019/Bilder/Evolution\\_1.jpg](https://cms.uni-konstanz.de/fileadmin/pi/fileserver/2019/Bilder/Evolution_1.jpg)

Caption: The Midas cichlid fish from the crater lakes of Nicaragua are one of the the best known examples for sympatric speciation. They evolved from a source population into a variety of independent species in less than 22,000 years.

<https://cms.uni-konstanz.de/fileadmin/pi/fileserver/2019/Bilder/Evolution.jpg>

Caption: Researchers found an increased activity of microRNA in young fish one day after they hatched.

Wissenschaftliche Ansprechpartner:

Prof. Axel Meyer, Dr. Paolo Franchini  
University of Konstanz

Originalpublikation:

Paolo Franchini, Peiwen Xiong, Carmelo Fruciano, Ralf F Schneider, Joost M Woltering, C Darrin Hulsey, Axel Meyer, MicroRNA gene regulation in extremely young and parallel adaptive radiations of crater lake cichlid fish, Molecular Biology and Evolution, msz168

Link: <https://academic.oup.com/mbe/advance-article/doi/10.1093/molbev/msz168/5545549>

Julia Wandt | idw - Informationsdienst Wissenschaft

**Further reports about:** > [Molecular Biology](#) > [cichlid fish](#)  
> [ecological niches](#) > [gene flow](#) > [genes](#) > [molecular mechanisms](#)  
> [new species](#) > [sympatric speciation](#)

**More articles from Life Sciences:**

Single enzyme helps drive inflammation in mice, provides target for new sepsis drugs

14.08.2019 | University of California - San Diego

Supercapacitors turbocharged by laxatives

13.08.2019 | University of Bristol

**All articles from Life Sciences >>>****The most recent press releases about innovation >>>****Die letzten 5 Focus-News des innovations-reports im Überblick:****Im Focus: Vehicle Emissions: New sensor technology to improve air quality in cities**

Researchers at TU Graz are working together with European partners on new possibilities of measuring vehicle emissions.


Today, air pollution is one of the biggest challenges facing European cities. As part of the Horizon 2020 research project CARES (City Air Remote Emission...

**Im Focus: Self healing robots that "feel pain"**

Over the next three years, researchers from the Vrije Universiteit Brussel, University of Cambridge, École Supérieure de Physique et de Chimie Industrielles de la ville de Paris (ESPCI-Paris) and Empa will be working together with the Dutch Polymer manufacturer SupraPolix on the next generation of robots: (soft) robots that 'feel pain' and heal themselves. The partners can count on 3 million Euro

in support from the European Commission.


Soon robots will not only be found in factories and laboratories, but will be assisting us in our immediate environment. They will help us in the household, to...

**NASA | A Year in the Life of Earth's CO2**

NASA Computer Model Provides a New Portrait of Carbon Dioxide

**Black Holes Come to the Big Screen**

The new movie "Interstellar" explores a longstanding fascination, but UA astrophysicists are using cutting-edge technology to go one better.

**NASA's Swift Mission Observes Mega Flares from a Mini Star**

NASA's Swift satellite detected the strongest, hottest, and longest-lasting sequence of stellar flares ever seen from a nearby red dwarf star.

**NASA | Global Hawks Soar into Storms**


NASA's airborne Hurricane and Severe Storm Sentinel or HS3 mission, will revisit the Atlantic Ocean for the third year in a row.

**Baffin Island - Disappearing ice caps**

Giff Miller, geologist and paleoclimatologist, is walking the margins of melting glaciers on Baffin Island, Nunavut, Canada.

**The sun's magnetic field is about to flip**


Something big is about to happen on the sun.

**The Infrasound Network and how it works**

The CTBTO uses infrasound stations to monitor the Earth mainly for atmospheric explosions.


**CU-Boulder team develops swarm of pingpong ball-sized robots**

Overview of more  
**VideoLinks >>>**

**Im Focus: Scientists create the world's thinnest gold**

Scientists at the University of Leeds have created a new form of gold which is just two atoms thick - the thinnest unsupported gold ever created.


The researchers measured the thickness of the gold to be 0.47 nanometres - that is one million times thinner than a human finger nail. The material is regarded...

**Im Focus: Study on attosecond timescale casts new light on electron dynamics in transition metals**

An international team of scientists involving the Max Planck Institute for the Structure and Dynamics of Matter (MPSD) in Hamburg has unraveled the light-induced electron-localization dynamics in transition metals at the attosecond timescale. The team investigated for the first time the many-body electron

dynamics in transition metals before thermalization sets in. Their work has now appeared in Nature Physics.

The researchers from ETH Zurich (Switzerland), the MPSD (Germany), the Center for Computational Sciences of University of Tsukuba (Japan) and the Center for...

**Im Focus: New approach in the fight against viruses**

In the ViroSens project, researchers from the Fraunhofer-Gesellschaft in Sulzbach and Regensburg are working together with industrial partners on a novel analytical method to make the potency testing of vaccines more efficient and cost-effective. The method combines electrochemical sensor technology and biotechnology and, for the first time, enables a completely automated analysis of the infection status

of test cells.

Most people see vaccinations as a benefit of modern medicine, as they protect against dangerous viral diseases. However, before a vaccine can enter the...

**All Focus news of the innovation-report >>>**

Top

Send this article

Print

Jetzt

12. 22. September

IAA

Home About us Partner Contact Sitemap find and help Deutsch Disclaimer

© 2000-2019 by innovations-report