

## Science News

from research organizations

### 'Left-handed' fish and asymmetrical brains

*Date:* November 15, 2017

*Source:* University of Konstanz

*Summary:* Biologists have discovered the relationship between “handedness”, brain structure and genes in extremely specialized cichlid fish.

*Share:* [f](#) [🐦](#) [G+](#) [p](#) [in](#) [✉](#)

#### FULL STORY



Cichlid fry with bent jaw.

*Credit: Image courtesy of University of Konstanz*

To humans, being right-handed or left-handed plays an important role. The majority favours the right side of their bodies, while only about three per cent of people innately use both hands equally well. Preferring one side of the body over another is not unique to humans: the

phenomenon occurs in the animal kingdom as well, for instance in great apes or birds, although their "handedness" is typically not as obvious as it is in humans. "Handedness" is particularly pronounced in cichlid fish of the species *Perissodus microlepis*, which is endemic to Lake Tanganyika in Africa.

This cichlid is highly specialised, living exclusively of the scales of other fish that they obtain from one side of the bodies of their prey only. When hunting, not only do individual fish display a pronounced predilection for attacking their prey from one direction; this preference is also mirrored by the shape of their heads: many individuals of this cichlid have asymmetrical jaws that bend to the left or right. Biologists from the University of Konstanz examined the relationship between the cichlids' preference for one side, the structure of their brains and their gene expression. A team led by Professor Axel Meyer, Professor of Zoology and Evolutionary Biology at the University of Konstanz, was able to demonstrate that a preference for one side in the animal's feeding behaviour correlates with a corresponding asymmetry of the cerebral hemispheres as well as unequal gene expression in the different hemispheres. The research results have been published in the current issue of the science journal *Genome Biology and Evolution*.

"Our research helps us to understand handedness and hemispheric asymmetries better. We hope that our results also will contribute to further our understanding of handedness in humans," explains the Konstanz biologist Dr Ralf Schneider, who, along with Dr Hyuk Je Lee, is one of the lead authors of the study.

In their study, the biologists documented the preference for one side of approximately 40 individual cichlids, followed by an examination of their brains. "We found a clear asymmetry across the animals' brains, especially in the Tectum Opticum region of the brain. However, this asymmetry was present only in individuals who displayed a distinct preference for one side when hunting," says Ralf Schneider. The Tectum Opticum is the region in the brain responsible for processing vision. In the respective cichlids, the hemisphere of the Tectum Opticum that controls the eye on the preferred side of the body was more strongly developed than the other. Individuals that did not display a marked preference for one side showed greater symmetry within the Tectum Opticum. The same proved true for other regions of the animals' brains. "This is interesting because it suggests a functional relationship between the shape of the brain and behavioural laterality," says Schneider.

Analyses of the genes (gene expression profiling) across several parts of the brain showed which genes tend to be expressed more strongly in the respective regions of the brain. The analyses revealed similar patterns to what the researchers had found in the fishes' brains structures: in the Tectum Opticum region alone, as much as 140 genes showed differing activity in the two hemispheres -- corresponding with the animals' preference for one side, gene activity in the respective hemispheres of the brain was increased or decreased. The most pronounced correspondence was determined for the gene synuclein gamma alpha (*sncga*) -- a particularly interesting result because previous studies on zebrafish have shown that this gene is expressed predominantly in the so-called habenula, a part of the brain that is asymmetric in all vertebrates, including humans. Besides synuclein gamma alpha, a range of other genes, some of which are associated with handedness and schizophrenia in humans, showed expression patterns in the brain hemispheres of the cichlids that suggest a link to handedness. There is reason to believe that some asymmetries in the brains of vertebrates could have originated in the habenula's asymmetry. The cichlids' preference for feeding on scales from just one side of their prey's body, as well as the asymmetrical shapes of their heads and brains, probably evolved due to the extremely fast "side-specific" development of gene regulation in their brains, the biologists suggest.

#### **Story Source:**

Materials

provided by **University of Konstanz**. *Note: Content may be edited for style and length.*

**Journal Reference:**

1. Hyuk Je Lee, Ralf F Schneider, Tereza Manousaki, Ji Hyoun Kang, Etienne Lein, Paolo Franchini, Axel Meyer. **Lateralized Feeding Behavior Comes with Asymmetrical Neuroanatomy and Lateralized Gene Expressions in the Brain in Scale-Eating Cichlid Fish.** *Genome Biology and Evolution*, 2017; DOI: 10.1093/gbe/evx218

**Cite This Page:**

MLA

APA

Chicago

University of Konstanz. "'Left-handed' fish and asymmetrical brains." ScienceDaily. ScienceDaily, 15 November 2017. <[www.sciencedaily.com/releases/2017/11/171115092045.htm](http://www.sciencedaily.com/releases/2017/11/171115092045.htm)>.

**RELATED STORIES****Baby Fish Exercising, a Surprising Source of Adaptive Variation in Fish Jaws**


Aug. 1, 2017 — A frustration of evolutionary biologists is that genetics can account for only a small percent of variation in physical traits. Now researchers have found new results on how another factor, a ... **read more »**

**Evolutionary Split Up Without Geographic Barriers**

July 5, 2016 — Evolutionary biologists have completed the most extensive study of sympatric speciation so far. They used around 20,000 characteristics of 450 fish to document the parallel evolution of cichlid fish ... **read more »**

**Study Finds Link Between Handedness, Mathematical Skills**

May 5, 2016 — The relationship between handedness and mathematical abilities is controversial. Now researchers say that a link between handedness and mathematical skills exists, but is more complex than is ... **read more »**

**Research Team Sheds Light on 'Rightie' or 'Leftie' Behavior in a Scale-Eating Cichlid**

Feb. 26, 2016 — Behavioral laterality, or left- or right-handedness, has been reported in many animals, including humans, chimpanzees, toads, rats, mice, and invertebrates such as crustaceans and insects. Now ... **read more »**

**FROM AROUND THE WEB**

*Below are relevant articles that may interest you. ScienceDaily shares links and proceeds with scholarly publications in the TrendMD network.*

**Persistence and the global dynamics of the positive solutions for a ratio-dependent predator-prey system with a crowding term in the prey equation**

Xianzhong, Acta Mathematica Scientia

## **Gathering Opinions on Depression Information Needs and Preferences: Samples and Opinions in Clinic Versus Web-Based Surveys**

Matthew T Bernstein et al., JMIR Publications

## **Peripheral blood RNA gene expression in children with pneumococcal meningitis: a prospective case-control study**

Benard W Kulohoma et al., BMJ Paediatrics Open

## **Survey of nutritional practices during therapeutic hypothermia for hypoxic-ischaemic encephalopathy**

Beth Hazeldine et al., BMJ Paediatrics Open

## **An acoustic Maxwell's fish-eye lens based on gradient-index**

Yuan, BG et al., Selection of Zero-Citation Articles: Science of Library and Information (ISTIC)

## **Failure laws of narrow pillar and asymmetric control technique of gob-side entry driving in island coal face**

Yang, International Journal of Mining Science and Technology

## **Child homicide perpetrators worldwide: a systematic review**

Heidi Stöckl et al., BMJ Paediatrics Open

## **Overexpression of Ribosomal RNA in the Development of Human Cervical**

Zhou, H et al., Selection of Zero-Citation Articles: Science of Library and Information (ISTIC)

---

Powered by

## **Free Subscriptions**

---

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

 Email Newsletters

 RSS Feeds

## **Follow Us**

---

Keep up to date with the latest news from ScienceDaily via social networks:

 Facebook

 Twitter

 Google+

 LinkedIn

## **Have Feedback?**

---

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

 [Leave Feedback](#)

 [Contact Us](#)

[About This Site](#) | [Editorial Staff](#) | [Awards & Reviews](#) | [Contribute](#) | [Advertise](#) | [Privacy Policy](#) | [Terms of Use](#)

Copyright 2017 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners. Content on this website is for information only. It is not intended to provide medical or other professional advice.

Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners.

Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.

