

## Science News

from research organizations

### Lungfish fins reveal how limbs evolved

*Date:* August 19, 2020

*Source:* University of Konstanz

*Summary:* New research on the fin development of the Australian lungfish elucidates how fins evolved into limbs with hands with digits. The main finding is that in lungfish a primitive hand is already present, but that functional fingers and toes only evolved in land animals due to changes in embryonic development.

*Share:* [!\[\]\(17413706fd4997a1a4bdf85c6864eee1\_img.jpg\)](#) [!\[\]\(f419710cbe076aa30a9c6c031b5cbe84\_img.jpg\)](#) [!\[\]\(2726020a4107bdc9042b257034f90eb3\_img.jpg\)](#) [!\[\]\(9459655bf14a84f4d775e8d814cca8c9\_img.jpg\)](#) [!\[\]\(de47dbdca34225b222a4a87ac0e499b3\_img.jpg\)](#)

#### FULL STORY

The evolution of limbs with functional digits from fish fins happened approximately 400 million years ago in the Devonian. This morphological transition allowed vertebrates to leave the water to conquer land and gave rise to all four-legged animals or tetrapods -- the evolutionary lineage that includes all amphibians, reptiles, birds and mammals (including humans). Since the nineteenth century several theories based on both fossils and embryos have been put forward trying to explain how this transformation unfolded. Yet, exactly how hands with digits originated from fish fins remained unknown.

An international team of biologists based at the University of Konstanz (Germany), Macquarie University in Sydney (Australia) and the Stazione Zoologica Anton Dohrn in Naples (Italy) has determined how limbs have evolved from fins using embryos of the Australian lungfish (*Neoceratodus forsteri*) for their study. The Australian lungfish is the closest living fish relative of tetrapods and is often considered a "living fossil" as it still resembles the fishes that were around at the time when the first four-limbed vertebrates began to walk on land. For these reasons the fins of lungfish provide a better reference to study the evolutionary transition of fins into limbs than any other extant fish species.

The team's research, which is reported in the latest issue of *Science Advances*, shows that a primitive hand is present in lungfish fins but at the same time suggests that the unique anatomy of limbs with digits only evolved during the rise of tetrapods through changes in embryonic development.

#### Insights from embryonic development: limb "architect" genes

To solve the puzzle of how limbs emerged from fins during evolution researchers have focused on embryonic development. "During embryogenesis, a suite of 'architect' genes shapes an amorphous group of precursor cells into fully grown limbs," explains Dr. Joost Woltering, first author on the study and an assistant professor in the Evolutionary Biology group at the University of Konstanz led by Professor Axel Meyer. The very same "architect" genes also drive fin development. However, because evolutionary changes have occurred in the activity of these genes, the developmental process produces fins in fish and limbs in tetrapods.

To compare this process in fins and limbs, the team studied such "architect" genes in the embryos of the Australian lungfish. "Amazingly, what we discovered is that the gene specifying the hand in limbs (*hoxa13*) is activated in a similar skeletal region in lungfish fins," explains Woltering. Importantly, this domain has never been observed in the fins of other fish that are more distantly related to tetrapods. "This finding clearly indicates that a primitive hand was already present in the ancestors of land animals."

## Developmental patterns: differences and similarities

The lungfish "hand," in spite of this modern genetic signature, only partially resembles the anatomy of tetrapod hands because it lacks fingers or toes. To understand the genetic basis for this difference the team went on to analyse additional genes known to be associated with the formation of digits, finding that one gene important for the formation of fingers and toes (*hoxd13* -- a "sister gene" to the above-mentioned *hoxa13*) appeared to be switched on differently in fins.

During tetrapod limb development, the *hoxd13* gene is switched on in a dynamic manner. It first becomes activated in the developing pinky finger and then expands all the way throughout the future hand towards the thumb. This process coordinates the correct formation of all five fingers. While Joost Woltering's team observed a similar activation pattern of this gene in lungfish fins, it did not show this expansion but only remained activated in exactly one half of the fin. Additional differences were found for genes that are normally switched off in digits. In lungfish fins these genes remain active, but on the opposite side of the domain where *hoxd13* is activated.

## Old hypotheses -- future directions

"All of this goes to show that while lungfish fins unexpectedly have a primitive hand in common with tetrapods, the fins of our ancestors also needed an evolutionary 'finishing touch' to produce limbs. In this sense it looks as if the hand was there first, only to be complemented with digits later during evolution," says Woltering. One influential hypothesis regarding the evolution of limbs first put forward by early 20th-century palaeontologists Thomas Westoll and William Gregory, and in the 1980s famously developed further by Neil Shubin, postulates that fingers and toes arose through an expansion of the skeletal elements on one side of the fins of the tetrapod ancestor. This inferred expansion of fin elements corresponds exactly to the differences the team found in the expansion of the digit genes between lungfish fins and tetrapod limbs. The team's observations on the activation and deactivation of limb "architect" genes in lungfish fins thus provides evidence in support of this classical transformational model.

In the future, to fully understand what causes this domain to expand, making our limbs so different from fish fins, the researchers plan to conduct further analyses on the development of fins and limbs, using lungfish but also more modern fish species such as cichlids as their embryos are easier to investigate using techniques like CRISPR. "To complete the picture of what happened in our fish ancestors that crawled onto land hundreds of millions of years ago, we really rely on currently living species to see how their embryos grow fins and limbs so differently," concludes Woltering.

## Background

- A new study by an international team of researchers from the University of Konstanz (Germany), Macquarie University in Sydney (Australia) and the Stazione Zoologica Anton Dohrn in Naples (Italy) provides an evolutionary model of how hands with digits emerged from fish fins.
- Studying the embryos of Australian lungfish (*Neoceratodus forsteri*), the closest extant fish relative of tetrapods, the researchers identified similarities and differences in the way lungfish fins and tetrapod limbs form during embryonic development.
- The presence of a primitive hand domain common to fins and limbs is revealed by the expression of a gene responsible for the specification of the hand in limbs (*hoxa13*). This gene becomes activated in similar skeletal domains in tetrapods and lungfish.
- One of the main morphological differences between fins and limbs, namely the absence of digits, can be explained by differences in the activation (*hoxd13*) and de-activation (*alx4*, *pax9*) of genes involved in digit development. This suggests that limbs with digits evolved from fish fins through changes in the activation of digit specific genes within a primitive hand domain.

---

## Story Source:

Materials provided by **University of Konstanz**. *Note: Content may be edited for style and length.*

---

## Journal Reference:

1. Joost M. Woltering, Iker Irisarri, Rolf Ericsson, Jean M. P. Joss, Paolo Sordino, Axel Meyer. **Sarcopterygian fin ontogeny elucidates the origin of hands with digits**. *Science Advances*, 2020; 6 (34): eabc3510 DOI: 10.1126/sciadv.abc3510
-

University of Konstanz. "Lungfish fins reveal how limbs evolved." ScienceDaily. ScienceDaily, 19 August 2020. <[www.sciencedaily.com/releases/2020/08/200819155723.htm](http://www.sciencedaily.com/releases/2020/08/200819155723.htm)>.

## RELATED STORIES

---

### Environmental Oxygen Triggers Loss of Webbed Digits

June 13, 2019 — Free fingers have many obvious advantages on land, such as in locomotion and grasping, while webbed fingers are typical of aquatic or gliding animals. But both amphibians and amniotes -- which ...

### Size Matters: To Livebearer Fish, Big Fins Are a Big Deal

Jan. 17, 2019 — Biologists studied the evolution of 40 molly and *Limia* species, and concluded dorsal fin displays arose first for males to compete with other males, only later being used in courtship displays to ...

### Researchers Pose Revolutionary Theory on Horse Evolution

Jan. 25, 2018 — While it is largely believed that horses simply evolved with fewer digits, researchers pose a new theory that suggests remnants of all five toes are still present within the hooves of the ...

### Vision, Not Limbs, Led Fish Onto Land 385 Million Years Ago

Mar. 7, 2017 — A new study suggests it was the power of the eyes and not the limbs that first led our aquatic ancestors to make the leap from water to land. The researchers discovered that eyes nearly tripled in ...

## FROM AROUND THE WEB

---

*Below are relevant articles that may interest you. ScienceDaily shares links with scholarly publications in the TrendMD network and earns revenue from third-party advertisers, where indicated.*

### Moving Along

Authors: Eugenio Guglielmelli et al., IEEE PULSE, 2015

### Retro Reproduction

Author: Shannon Fischer, IEEE PULSE, 2015

### Veblen's evolutionary economics of religion and the evolutionary psychology of religion

Charles G. Leathers et al., International Journal of Social Economics, 2014

### ESRA19-0195 Hand trauma: 'gradient of pressure' technique to block the radial and medial nerves

K Geier, Regional Anesthesia & Pain Medicine, 2019

### The Genie Is Out of the Bottle

Author: Colin J. H. Brenan, IEEE PULSE, 2015

### Comparison of methods of measurement of the finger flexor muscles' strength through dynamometry and modified manual sphygmomanometer

Lucareli, Paulo Roberto Garcia et al., Einstein (São Paulo), 2010

### Restoring the Sense of Touch: From "Sci-Fi Dream" to Reality

Author: Leslie Mertz, IEEE PULSE, 2020

### State of the ART

Author: Shannon Fischer, IEEE PULSE, 2015

I consent to the use of Google Analytics and related cookies across the TrendMD network (widget, website, blog).

Yes


No

## Free Subscriptions

---

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

 [Email Newsletters](#)

 [RSS Feeds](#)

## Follow Us

---

Keep up to date with the latest news from ScienceDaily via social networks:

 [Facebook](#)

 [Twitter](#)

 [LinkedIn](#)

## Have Feedback?

---

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

 [Leave Feedback](#)

 [Contact Us](#)

[About This Site](#) | [Staff](#) | [Reviews](#) | [Contribute](#) | [Advertise](#) | [Privacy Policy](#) | [Editorial Policy](#) | [Terms of Use](#)

Copyright 2020 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners.

Content on this website is for information only. It is not intended to provide medical or other professional advice.

Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners.

Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.

— [GDPR: Privacy Settings](#) —