This website uses cookies to ensure you get the best experience on our website. More info

Nanotechnology

Physics

Earth Astronomy & Space Technology

Chemistry

Biology Other Sciences

search

Biology

Evolution October 25, 2018

Team discovers the genetic basis for the repeated evolution of color patterns

October 25, 2018, University of Konstanz



Pundamilia nyererei fish from Lake Victoria that has been altered using CRISPR-Cas as 'gene scissors'. The fish demonstrates horizontal stripes on its flank as a result of the genetic alteration. Credit: Claudius Kratochwil

A team of evolutionary biologists from the University of Konstanz, headed by Prof. Dr. Axel Meyer, has discovered the genetic basis for the repeated evolution of colour patterns. The findings about the stripes of the diverse species of East African cichlid fishes explain how evolution can repeat itself at record speed. The study is published in Science magazine on 26 October.



Understand CRISPR

Optimize your gene-editing knowledge online from Harvard VPAL experts

Harvard's VPAL


OPEN >


Why does evolution repeat itself? What happens genetically during these repetitions? Do the same or other genes and mechanisms produce organisms that look similar? Professor Axel Meyer and his team from Konstanz University have come closer to answering this question, which is as old as it is important. The answer is quite astonishing. They studied a special colour pattern that is omnipresent among all kinds of animals: horizontal stripes. The researchers

were able to identify the basis of the repeated evolution of these stripes with modern methods in genomics and molecular biology, and verified them with a CRISPR-Cas mutant fish that added a stripe.

More than 1200 types of colourful cichlid can be found in the large African lakes Malawi, Victoria and Tanganjika. Not only are they diverse in colour, they also have numerous colour patterns such as horizontal or vertical stripes.







Subscribe

email

"But that's not all," explains Axel Meyer. "Cichlids are prime examples of evolution. They are extremely diverse in terms of social behaviour, body shape, colour pattern and many other biological aspects, but at the same time, certain features repeatedly evolved independently between the lakes." This principle of repeated evolution—biologists term it convergent evolution—makes cichlids the perfect target to study the genetic basis of this phenomenon. If similar colours and body shapes have emerged in several evolutionary lines independently from each other, this means that evolution reacted to similar environmental conditions in the same way. But how does such evolutionary repetition work genetically?



Colourful cichlids with horizontal stripes living in the large African lakes Malawi, Victoria and Tanganjika illustrate the repeating (convergent) process of evolution. Credit: Claudius Kratochwil

Which gene and which genetic mechanism are responsible for the cichlids' stripes to come and go has now been reconstructed in the laboratory through genome analyses, breeding and experiments, including CRISPR-Cas technology. Dr. Claudius Kratochwil, early career researcher in Professor Meyers' team and first author of the study in Science, explains: "In breeding experiments, we can exactly determine on which of the 22 chromosomes, even on which area of the chromosome of the fish, the genetic instruction for stripes is located."

Understand CRISPR

Optimize your gene-editing knowledge online from Harvard VPAL experts as.harvardx.harvard.edu

OPEN

The relevant gene on this chromosome part is called agrp2. This "stripe gene," its origin and distribution in other African lakes was described in comparative molecular studies. From an evolutionary point of view, the cichlids' stripes are unstable. Over the course of a few million years, they have been lost and re-emerged in the African lakes many times over. As these species (with and without stripes) are so young, they can be interbred in aquaria.

Breeding and examining the cichlids with and without stripes in the laboratory shows that all cichlids carry the "stripes gene," but the switches (regulatory elements) of this gene differ. This genetic switch causes the gene in species without stripes to be more activated. As a result, a lot of protein is produced. The 'stripes gene' agrp2 works as a 'stripes inhibitor': if gene production is high, stripes will be suppressed, if production is low, they will remain.

The researchers were able to demonstrate this by using modern genetic methods. "If we use CRISPR-Cas to remove the gene from the genome of a species without stripes," Kratochwil explains, "then even a 'stripeless' fish will

Relevant PhysicsForums posts

Chemical Computers 5 hours ago

could our nose and lungs filter tiny paint pieces 18 hours ago

Any bio-related benefit of UV radiation? Oct 27, 2018

What are the best references for neurogenesis? Oct 24, 2018

How does chromosome number get halved in meiosis I? Oct 24, 2018

Selection versus genetic drift Oct 22, 2018

More from Biology and Medical

suddenly develop stripes, as we showed with a CRISPR-Cas mutant fish. This proves that the stripes gene is the decisive genetic factor."



The especially colourful Haplochromis chilotes fish from Lake Victoria exhibits both horizontal and vertical stripes. Credit: Claudius Kratochwil

The latest findings on this genetic mechanism, the activation or deactivation of stripes by the "stripes gene," were published in the current issue of Science. Interestingly, the cichlid's agrp2 gene is a copy of the agouti gene in mammals, which is responsible for the different coat colours of cats, dogs, horses and striped baby birds.

"The world of animals might be much less colourful without the agouti gene family," says Claudius Kratochwil. The mechanism of the "stripes gene" in cichlids clearly makes repeated evolution possible within the briefest of times, relatively speaking. If characteristics are lost during evolution, usually this loss is permanent, as the Belgian palaeontologist Louis Dollo realized exactly 125 years ago, and wrote up his conclusions in "Dollos Law" in 1893. The special aspect of the stripes gene agrp2 is that it makes repeated evolution of a characteristic possible in a simple way. If a cichlid loses its stripes, that does not mean they will never return or vice versa. These molecular-biological studies also show that palaeontological rules and evolutionary rules have to be questioned once again.

Explore further: How the clownfish earned its stripes: Color pattern evolution in coral reef fishes

More information: Agouti-related peptide 2 facilitates convergent evolution of stripe patterns across cichlid fish radiations, Science (2018). science.sciencemag.org/cgi/doi ... 1126/science.aao6809

Journal reference: Science

O shares

Provided by: University of Konstanz

feedback to editors

66

G+

Tweet

Favorites



Understand CRISPR

Optimize your gene-editing knowledge online from Harvard VPAL experts

Related Stories



How the clownfish earned its stripes: Color pattern evolution in coral reef fishes September 4, 2018

Coral reef fishes, including clownfish, display a wide variety of colors but it remains unclear how these colors evolved or

published in BMC Biology sheds new light on the ...

Recommended for you



You are the company you keep—A new screening method detects direct biomolecule interactions October 26, 2018

Proteins are the building blocks of the cell. They do most of the work and are essential for the structure, function and dynamic regulation of the cell and body's tissues and organs.

Proteins rarely work alone, they interact, ...



Study finds Nemo's bright colours protect it from predators July 25, 2018

how they develop throughout a fish's life. Research

The colourful stripes of coral reef-dwelling clownfish may serve to warn predators about their poisonous anemone hosts, according to a new study by The University of Western Australia.



Animal migration requires both movement corridors and food October 26,

An analysis of mule deer migration routes in two areas of Wyoming documents the idea that the animals choose readily traversable pathways where they also can find food.

User comments

Please sign in to add a comment. Registration is free, and takes less than a minute. Read more

| email | |
|----------|--|
| password | |
| | |
| Sign in | |

Click here to reset your password.

Sign in to get notified via email when new comments are made.

We recommend

Promoter of IncRNA Gene PVT1 Is a Tumor-Suppressor DNA Boundary Element Cell

CRISPR-Cas9 screens in human cells and primary neurons identify modifiers of C9ORF72 dipeptiderepeat-protein toxicity

Nature Genetics

Comparative genome and methylome analysis reveals restriction/modification system diversity in the gut commensal Bifidobacterium breve

Nucleic Acids Research

Broad Institute Team Engineers New Dx Technology Powered by CRISPR, Isothermal Amplification 360Dx

Android app

Тор

Empire Genomics Inks Italian Distribution Deal 360Dx

Help

Broad Institute Team Engineers Sherlock, Powered by CRISPR, Isothermal Amplification 360Dx

Powered by TREND MD

I consent to the use of Google Analytics and related cookies across the TrendMD network (widget, website, blog). Learn more

Feature Stories

Yes

No

Connect

Home FAQ Sponsored Account Latest news iOS app

Science X Account

Search About Newsletter Week's top Amazon Kindle

Mobile version Contact RSS feeds Archive Push notification

© Phys.org 2003 - 2018, Science X network

Privacy Policy Terms of Use