

Your source for the latest research news

Science News

from research organizations

Rapid evolution: New findings on its molecular mechanisms

Date: August 14, 2019

Source: University of Konstanz

Summary: Evolutionary biologists have analyzed the role of microRNAs in the evolution of new species.

Share: $f \not y p$ in \square

FULL STORY

The mechanisms by which new species arise are still not fully understood. What are the evolutionary processes that drive the evolution of new species? Evolutionary biologists traditionally assumed that geographical barriers between animal populations play a decisive role (allopatric speciation): a species is physically separated into two or more isolated populations, thereby preventing gene flow between these groups. The subpopulations adapt to their respective habitats and evolve into independent species with different characteristics. In recent years, however, the evolutionary biologist Professor Axel Meyer from the University of Konstanz has not only been able to show that new species can evolve from a source population within a shared habitat and in the presence of gene flow (sympatric speciation), but that this type of speciation might be much more common than previously thought. His laboratory is investigating both the ecological and genetic mechanisms that facilitate sympatric speciation. In a recent publication in the scientific journal Molecular Biology and Evol lution, Axel Meyer and his colleagues Paolo Franchini, Peiwen Xiong, Carmelo Fruciano, Ralf Schneider, Joost Woltering and Darrin Hulsey identify the decisive role that a kind of genetic switch, microRNAs, play in sympatric speciation.

Rapid evolution of cichlids

The researchers led by Axel Meyer analyze the Midas cichlid fish from Nicaragua's volcanic crater lakes as a model system in their research. Cichlids are known for their ability to adapt to new environments at an exceptionally rapid rate and to form new species. The new fish species originate from a population found in the large lakes of Nicaragua, but after colonizing several very small and young crater lakes, they adapted to new ecological niches, evolved new characteristics (e.g. a more elongated body or a different jaw) and evolved into a variety of new

species in less than 22,000 years. A further distinctive feature of these Midas cichlids is that they evolved into different species within the same population, sometimes repeatedly, by adapting to different ecological niches within their small crater lakes. The Midas cichlids of Nicaragua are thus one of the best known examples for sympatric speciation. Within such a short time period new mutations are very unlikely to happen. This made finding a molecular mechanism that can bring about different body shapes, adaptations and thereby distinct ecological niches so difficult and interesting. The evolution of microRNAs and new target sites for them to regulate the expression of genes offers one potential molecular mechanism that can bring about very fast evolutionary change with very little genetic differences between the extremely young species.

microRNA

The biologists from Konstanz carried out genetic analyses of five species of Midas cichlids from the crater lakes Apoyo and Xiloá. Here, they focused in particular on the function of the so-called microRNA -- a non-coding ribonucleic acid that has a regulatory effect on gene expression. The researchers found an increased activity of microRNA in young fish one day after they hatched during a phase in which the fish bodies are formed. They analyzed the interaction between microRNA and gene expression, identifying specific pairs of microRNAs and genes that influence each other. The microRNA suppresses the expression of the target genes and thus has a regulatory effect on them: the more active a particular microRNA is, the more effectively the target gene is suppressed or "switched off." "Our research results provide strong evidence that extremely quickly evolving microRNA regulation contributes to the rapid sympatric speciation of Midas cichlids," says Paolo Franchini.

Story Source:

Materials provided by **University of Konstanz**. Note: Content may be edited for style and length.

Journal Reference:

1. Paolo Franchini, Peiwen Xiong, Carmelo Fruciano, Ralf F Schneider, Joost M Woltering, C Darrin Hulsey, Axel Meyer. **MicroRNA gene regulation in extremely young and parallel adaptive radiations of crater lake cichlid fish**. Molecular Biology and Evolution, 2019; DOI: 10.1093/molbev/msz168

Cite This Page:	MLA	APA	Chicago
-----------------	-----	-----	---------

University of Konstanz. "Rapid evolution: New findings on its molecular mechanisms." ScienceDaily. ScienceDaily, 14 August 2019. www.sciencedaily.com/releases/2019/08/190814105222.htm.

RELATED STORIES

Biologists Watch Speciation in a Laboratory Flask

Nov. 29, 2016 — The evolution of a new species can occur rapidly enough for them to observe the process in a simple laboratory flask, biologists have discovered. In a month-long experiment using a virus harmless to ... read more »

On the Origin of Eukaryotes: When Cells Got Complex

Feb. 3, 2016 — Just as physicists comprehend the origin of the universe by observing the stars and archeologists reconstruct ancient civilizations with the artifacts found today, evolutionary biologists study the ... **read more** »

How Anti-Evolution Bills Evolve

Dec. 17, 2015 — An evolutionary biologist has analyzed political opposition to evolution and found it has evolved. The researcher analyzed the text in anti-evolution legislation using software for building genetic ... **read more** »

Evolutionary Biologists Observe Parallel, Repeated Evolution of Cichlid Fish in Nicaragua

Oct. 28, 2014 — If one would rewind the tape of life, would evolution result in the same outcome? An evolutionary biologist came up with this famous question. He suggested that evolution would not repeat itself: the ... **read more** »

FROM AROUND THE WEB

Below are relevant articles that may interest you. ScienceDaily shares links with scholarly publications in the Trend-MD network and earns revenue from third-party advertisers, where indicated.

Genome-wide association study identifies loci for body shape in the large yellow croaker (Larimichthys crocea)

Dong et al., Aquaculture and Fisheries, 2019

Who eats whom, when and why? Juvenile cannibalism in fish Asian seabass

Liu et al., Aquaculture and Fisheries, 2017

Natural and artificial polyploids in aquaculture

Zhou et al., Aquaculture and Fisheries, 2017

Influence of sampling frequency on detectability of fish community and fish species in a fishery-independent survey

Zhao et al., Aquaculture and Fisheries, 2017

Sex-biased gene discovery from the gonadal transcriptomes of the large yellow croaker (Larimichthys crocea)

Xiao et al., Aquaculture and Fisheries, 2019

Phylogenomic Mining of the Mints Reveals Multiple Mechanisms Contributing to the Evolution of Chemical Diversity in Lamiaceae

Benoît Boachon et al., Molecular Plant, 2018

The Tea Tree Genome Provides Insights into Tea Flavor and Independent Evolution of Caffeine Biosynthesis

En-Hua Xia et al., Molecular Plant, 2017

Crop Breeding Chips and Genotyping Platforms: Progress, Challenges, and Perspectives

Awais Rasheed et al., Molecular Plant, 2017

Powered by TREND MD

I consent to the use of Google Analytics and related cookies across the TrendMD network (widget, website, blog). Learn more

Yes No

Free Subscriptions

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

- **f** Facebook
- in LinkedIn

Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

- Leave Feedback
- Contact Us

About This Site | Staff | Reviews | Contribute | Advertise | Privacy Policy | Editorial Policy | Terms of Use

Copyright 2019 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners. Content on this website is for information only. It is not intended to provide medical or other professional advice. Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners. Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.